

Available online at www.ijit.net**International Journal of Integrative sciences, Innovation and Technology (IJIT)**

(A Peer Review E-3 Journal of Science Innovation Technology)

Journal homepage: <http://www.ijit.net/>

eISSN 2278-1145

Research Unlimited

Vol. IV Iss 4

Research Paper

Microbial Quality of Fresh Lettuce Irrigated with Untreated Waste Water Around Aksum University, Central Zone of Tigray Region.

Teklemichael Tesfay^{a*}, Askalu Welearegay^b, Baymot Dufera^b, Betelihem Deribachew^b and Doboal Duany^b

^aDepartment of Biology, College of Natural and Computational Sciences, Aksum University, Aksum, Ethiopia, P.O. Box: 1010, N E Africa

^bDepartment of Biotechnology, College of Natural and Computational Sciences, Aksum University, Aksum, Ethiopia, P.O. Box: 1010, N E Africa.

ARTICLE INFO

Article history:

Received 17 July 2015

Received in revised form 21 July 2015

Accepted 07 August 2015

Keywords:

Coliforms

Lettuce

Salmonella spp.

Waste water

ABSTRACT

Vegetables are essential components of the human diet and there is considerable evidence of the health and nutritional benefits related with the consumption of fresh vegetables. However, they can cause health problem if contaminated by pathogenic bacteria. The present study focused on the microbial quality of vegetables irrigated with untreated waste water. Data were analyzed using percentage and tables. The result of the study shows that 100% of the lettuce samples and 75% of the water samples were positive for salmonella spp. On the other hand, higher numbers of coliforms from lettuce were counted ranging from 3.207×10^3 - 4.063×10^5 cfu/ml. Many people consume and sold lettuce in nearby towns and cause major health problems. Therefore, the present study can be used as an alarm for the concerned bodies to take measure.

© 2012 Editor-IJIT. Hosting by AGSI Publications. All rights reserved.

How to cite this article: Teklemichael Tesfay, Askalu Welearegay, Baymot Dufera, Betelihem Deribachew and Doboal Duany (2015). Microbial Quality of Fresh Lettuce Irrigated with Untreated Waste Water Around Aksum University, Central Zone of Tigray Region, International Journal of integrative Sciences, Innovation and Technology (IJIT), 4(4), 4 – 6.

1. Introduction

Fresh fruit and vegetables are important components of the human diet and there is considerable evidence of the health and nutritional benefits associated with their consumption. In the USA, Canada, New Zealand and several European Union states, public health institutions have run campaigns recommending the daily consumption of at least five daily servings of fruit and vegetables. As well as a significant rise in the consumption of fresh produce for health benefits, there have also been significant changes in lifestyles and major shifts in consumption trends¹.


Vegetable salad is a very common food accompaniment in many countries. The vegetables that usually make up this method include tomatoes, cucumber, carrots, green chili, cabbage and lettuce. They are sold in almost every market, and can be seen hawked around by traders. Fruits and vegetables have been identified as significant sources of pathogens and chemical contaminants².

Vegetables have been associated with outbreaks of food borne disease in many countries. Organisms involved include bacteria, viruses and

* Corresponding author. Tel.: +0-000-000-0000 ; fax: +0-000-000-0000.

E-mail address: teklemichael2010@gmail.com

Peer review under responsibility of board members IJIT.



IJIT/ – see front matter ©2012editor.ijit. Hosting by AGSI Publications. All rights reserved.
<http://ijit.net>

parasites. Raw vegetables can harbour many microorganisms, which may be dispersed over the plant or appear as microcolonies fixed in the plant tissue³. The majority of microorganisms associated with raw vegetables are gram negative organisms tend to dominate the microbial population. Vegetables are highly exposed to microbial contamination through contact with soil, dust and water and by handling at harvest or during postharvest processing. They therefore harbour a different type of microorganisms including plant and human pathogens⁴.

Differences in microbial profiles of various vegetables result largely from unrelated factors such as resident microflora in the soil, application of nonresident micro flora via animal manures, sewage or irrigation water, transportation and handling by individual retailers⁵. Use of untreated waste water and manure as fertilizers for the production of fruits and vegetables is the main contributing factor to contamination⁶. Tomato, Cucumber, Carrot, Green chili, Lemon, Coriander leaf, Pepper mint and Beet root are some of the vegetables that are normally consumed raw in order to obtain their valuable nutrients in best form and their traditional use in best form and their traditional use in preparing salads is familiar throughout the world in the same manner.

Commercial and small-scale farmers generally irrigate their produce with water from nearby rivers, streams, ponds and dams most of which do not meet the required standard for irrigation⁷. The microbiological quality of irrigation water is critical because water contaminated with animal or human wastes can introduce pathogens during pre-harvesting and post-harvest⁸. Furthermore, ⁹ reported that contaminated irrigation water and surface run-off water might be major source of pathogenic microorganisms that contaminate fruits and vegetables in fields. Food safety issues are of growing concern to consumers globally because of the risk associated with consumption of foods contaminated with pathogens in irrigated vegetables. In the study area farmers were commonly cultivated vegetables irrigated with waste water drained from different cafeterias. Therefore, the objective of the current study is to assess the quality of lettuce irrigated with waste water around Aksum University.

2. Materials and Methods

2.1 Sample size and Sampling Techniques

The study was conducted for about 3 months from April to June 2014. A total of 8 samples of lettuce and waste water were collected randomly at different times. The raw lettuce was collected by using sterile glove and placed in sterile polyethylene plastic bags while the waste water samples were placed in sterile bottles. Samples were taken to the laboratory and processed immediately within 2 hours.

2.2 Isolation Techniques of salmonella spp.

A 25 gram of lettuce sample were weighed and immersed in 250 ml beaker containing 100 ml of sterile distilled water. From the rinsed water a serial dilutions were prepared. Media were prepared according to the manufacturer's guide. Finally, a 0.1 ml of each dilution was spread plated on SS-agar and incubated at 37⁰c for 24 hours. Finally, colonies with black centers were confirmed by using, triple sugar iron agar test, motility test and colors of the colony¹⁰.

2.3 Coliform count using MacConkey agar

The enumeration of coliforms was done using MacConkey agar following the methods recommended by¹¹. Media were prepared according to the manufacturer's guide. A 25 gram of lettuce sample

were weighed and immersed in 250 ml beaker containing 100 ml of sterile distilled water. From the rinsed water a serial dilution were prepared. Volumes of 1 ml of appropriate dilutions were pour plated to MacConkey agar and plates were incubated at 37⁰C for 24 hours. Finally, colonies between 30-300 were counted.

3. Result and Discussion

Table 1. Isolation of Salmonella spp. from lettuce and Waste Water

Sample source	Number of samples	Salmonella species	
		Positive	Negative
Lettuce	4	4(100%)	-
Waste water	4	3(75%)	1 (25%)

According to the above Table, 75% of the lettuce samples were positive for *salmonella* pp. The current result agreed with the works of ¹² where they detect *Salmonella* spp. in 80% of waste water samples. Only 25% of the water samples were free of *salmonella*. Our study also shows that 100% of *Salmonella* spp. were detected in raw lettuce samples. This is in line with the results of ¹³ where they isolate *Salmonella* spp. in 20% of raw lettuce samples. According to ¹⁴ guide lines in 25 gram of raw vegetables *Salmonella* spp. should not be detected. However, in most of the samples *salmonella* spp. were detected. The existence of *salmonella* spp. in lettuce is due to the use of waste water for irrigation purpose. As it was observed from the preliminary survey, farmers use waste water for irrigation purpose to cultivate vegetables. This can result in major health problem of the society if care is not taken by the consumers. *Salmonella* spp. were isolated from waste water, indicating that waste water is the major source of contamination for lettuce in the agricultural field. The source of *Salmonella* to the waste water could be the student cafeterias where utensils used for the preparation of food were washed every day and the left over of the food may be discharged to the river. According to ¹⁵, effect of microbiological hazards such as *Salmonella* on food safety is now a major public health concern worldwide. However, our result shows that all of the samples were positive for *salmonella* spp.

Table 2. Coliform count from lettuce sample

Sample source (lettuce)	Cfu/ml
sample 1	4.063X10 ⁵
sample 2	5.031X10 ³
Sample 3	3.207X10 ³
Sample 4	4.873X10 ³

The above Table 2 shows coliform count ranging from 3.207X10³-4.063X10⁵. Our current study is similar with the works of ¹⁶ where they reported coliform counts of 3.7 ± 0.5 cfu/g in lettuce samples. According to ¹⁴, the accepted value for coliform count in fruits and fresh vegetables should be <10⁴ cfu/ml. However, our current result exceeds this value. This indicates that the lettuce which is cultivated by the farmers is poor in quality. The higher number of coliforms indicates that pathogenic bacteria may exist in the lettuce.

Table 3. Coliform count from waste water

Sample source (waste water)	Cfu/ml
sample 1	2.54 X10 ⁵
sample 2	1.555 X10 ⁵
Sample 3	3.754 X10 ⁴

Sample 4	3.163 X10 ⁴
----------	------------------------

The above Table indicates that the waste water which is used for irrigation purpose contains higher number of coliforms. The number of colony forming units (cfu/ml) of the sample ranges from 3.163 X10⁴-2.54 X10⁵. Farmers use this water for irrigation purpose without any treatment. As a result, freshly consumed fruits and vegetables may be easily contaminated by pathogenic bacteria. The higher numbers of coliforms in food samples indicate that the food may contain pathogenic bacteria. This was confirmed by detecting *salmonella* spp. from both lettuce and waste water. The existence of *Salmonella* spp. in the water samples may be from left over foods that comes from student cafeteria.

4. Research Highlights

In the current study pathogenic bacteria (*Salmonella* spp.) were detected in both waste water used for irrigation purpose and the lettuce.

5. Limitations

Due to limitation of some media in the laboratory no other pathogenic bacteria were detected.

6. Recommendations

- Vegetables cultivated using waste water should not be consumed
- The Ethiopian government should have standards for microbial quality of fresh vegetables

7. Conclusion

In conclusion, the high coliform load in lettuce samples could serve as an indicator for the need to promote awareness about the possible health hazards that could be due to poor handling of these vegetables such as lettuce. On the other hand, the detection of *Salmonella* spp. in the lettuce samples could cause a major risk in the society. Therefore, the need for regulatory bodies to ensure that microbiological standards are established and practiced by farmers and marketers for the handling and distribution of vegetables is required.

Acknowledgements

We would like to thank Aksum University for supporting all the materials and chemicals required for the research work.

REFERENCES

1. Vose, D., Koupeev, T. and Mintiens, K. (2010). A Quantitative Microbiological Risk Assessment of *Salmonella* spp. in broiler (*Gallus gallus*) meat production. Technical Report submitted to EFSA. <http://www.efsa.europa.eu/en/supporting/pub/183e.htm>.
2. Uzeh, R., Alade, F. and Bankole, M. (2009). The microbial quality of prepacked mixed vegetable salad in some retail outlets in Lagos, Nigeria. *Afr. J. Food Sci.* **3** (9), 270-272.
3. Szabo, E.A. and Coventry, M.J. (2001). Vegetables and vegetable products. In C.J. Moir, et al., (eds) Spoilage of Processed Foods; Causes and Diagnosis. AIFST Inc. (NSW Branch) Food Microbiology Group. Waterloo.
4. Nguyen, C. and Carlin, F. (1994). The microbiology of minimally processed fresh fruits and vegetables. *Crit. Rev. Food Sci. Nutr.*, **34**, 371- 401.
5. Ray, B. and Bhunia, A.K. (2007). *Fundamental Food Microbiology*. 4th edn. CRC Press, USA, 2007.
6. Olayemi, A. (1997). Microbiological hazards associated with agricultural utilization of urban polluted river water. *International J. Enviro. Health Res.*, **7**(2), 149-154.
7. Westcot, D. W. (1997). Quality control of wastewater for irrigated crop production. FAO. Water Reports n^o10. Rome.
8. FDA, Safe Handling of Raw Produce and Fresh-Squeezed Fruit and Vegetable Juices. United States Department of Health and Human Services, Food and Drug Administration, Center for Food Safety and Applied Nutrition, 2005.
9. Palese, A., Pasquale, V., Celano, G., Figliuolo, G., Masi, S. and Xiloyannis, C. (2009). Irrigation of olive groves in Southern Italy with treated municipal wastewater: Effects on microbiological quality of soil and fruits agriculture. *Ecosystems and Environment*, **129**, 43-51.
10. Khan, M., Saha, M. and Kibria, A. (1992). A bacteriological profile of salad vegetables in Bangladesh with special reference to coliforms. *Applied Microbiology* **14**, 88-90.
11. Roberts, D. and Greenwood, M. (2003). *Practical food microbiology* 3rd edn. Blackwell publishing Ltd.
12. Melloul, A., Hassani, L. and Rafouk, L. (2001). *Salmonella* contamination of vegetables irrigated with untreated waste water. *World Journal of Microbiology and Biotechnology* **17**, 207-209.
13. Itohan, M., Peters, A. and Kolo, I. (2011). Bacterial contaminants of salad vegetables in Abuja Municipal Area Council, Nigeria. *Malaysian Journal of Microbiology*, **7**(2), 111-114.
14. PHLS (2000). Guidelines for the microbiological quality of some ready to eat foods at the point of sale.
15. WHO (2002). Global Strategy for Food Safety: Safer Food for Better Health. Food Safety Issues, WHO Geneva.
16. Cobbina, J., Kotochi, M., Korese, J. and Akrong, M. (2013). Microbial Contamination in Vegetables at the Farm Gate Due to Irrigation with Wastewater in the Tamale Metropolis of Northern Ghana. *Journal of Environmental Protection*, **4**, 676-682.